



YOUR GUIDE TO TREATMENT WITH

TIBSOVO[®]

ivosidenib tablets 250mg

WHAT IS TIBSOVO[®]?

TIBSOVO is a prescription medicine used to treat adults with bile duct cancer (cholangiocarcinoma) that has spread:

- who have already received previous treatment(s) and
- whose tumor has a certain type of abnormal isocitrate dehydrogenase-1 (IDH1) mutation.

Your healthcare provider will perform a test to make sure that TIBSOVO is right for you.

It is not known if TIBSOVO is safe and effective in children.

WHAT ARE THE POSSIBLE SIDE EFFECTS OF TIBSOVO?

TIBSOVO may cause serious side effects, including QTc prolongation and Guillain-Barré syndrome.

Please see additional Important Safety Information throughout and accompanying Full Prescribing Information, including Medication Guide.

BEFORE TAKING TIBSOVO®, TELL YOUR HEALTHCARE PROVIDER ABOUT ALL OF YOUR MEDICAL CONDITIONS, INCLUDING IF YOU:

- Have any heart problems, including a condition called long QT syndrome
- Have problems with abnormal electrolytes, such as sodium, potassium, calcium, or magnesium levels
- Have nervous system problems
- Have problems with your kidneys or are on dialysis
- Have any liver disorders, including cirrhosis
- Are pregnant or plan to become pregnant. TIBSOVO may cause harm to your unborn baby. You should avoid becoming pregnant during treatment with TIBSOVO. Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with TIBSOVO
- Are breastfeeding or plan to breastfeed. It is not known if TIBSOVO passes into your breast milk. Do not breastfeed during your treatment with TIBSOVO and for at least 1 month after your last dose of TIBSOVO

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your healthcare provider if you take hormonal contraceptives.

TIBSOVO may affect how hormonal contraceptives work and may cause them to not work as well.

HOW SHOULD I TAKE TIBSOVO?

- Take TIBSOVO exactly as your healthcare provider tells you to
- Do not change your dose or stop taking TIBSOVO without talking to your healthcare provider
- Take TIBSOVO 1 time a day about the same time each day
- Swallow TIBSOVO tablets whole. Do not split, crush, or chew the tablet
- TIBSOVO can be taken with or without food
- Do not take TIBSOVO with a high-fat meal. An example of a high-fat meal includes 2 eggs fried in butter, 2 strips of bacon, 2 slices of white bread with butter, 1 croissant with 1 slice of cheese, and 8 ounces of whole milk (approximately 1,000 calories and 58 grams of fat)
- If you vomit after taking a dose of TIBSOVO, do not take an additional dose. Take your next dose at your usual time
- If you miss a dose of TIBSOVO or did not take it at the usual time, take your dose as soon as possible and at least 12 hours before your next dose. Return to your normal schedule the following day. **Do not** take 2 doses of TIBSOVO within 12 hours

Please see accompanying Full Prescribing Information, including Medication Guide.

 **TIBSOVO®**
ivosidenib tablets 250mg

WHAT ARE THE POSSIBLE SIDE EFFECTS OF TIBSOVO®?

TIBSOVO may cause serious side effects, including:

- **Changes in the electrical activity of your heart called QTc prolongation. QTc prolongation can cause irregular heartbeats that can be life-threatening.** Your healthcare provider will check the electrical activity of your heart with a test called an electrocardiogram (ECG) during treatment with TIBSOVO. Tell your healthcare provider right away if you feel dizzy, lightheaded, or faint
- **Guillain-Barré syndrome** has happened in people treated with TIBSOVO. Your healthcare provider will monitor you for nervous system problems and will permanently stop your treatment with TIBSOVO if you develop Guillain-Barré syndrome. Tell your healthcare provider right away if you develop any signs or symptoms of Guillain-Barré syndrome, including:
 - Weakness or tingling feeling in your legs, arms, or upper body
 - Any changes in your ability to see, touch, hear, or taste
 - Numbness and pain on one side or both sides of your body
 - Burning or prickling sensation
 - Difficulty breathing

The most common side effects of TIBSOVO in adults with cholangiocarcinoma include:

- Fatigue
- Nausea
- Abdominal pain
- Diarrhea
- Cough
- Decreased appetite
- Fluid and swelling in your stomach area
- Vomiting
- Hemoglobin decreased (anemia)
- Rash
- Changes in liver function tests

Your healthcare provider will do blood tests before you start and during treatment with TIBSOVO. Your healthcare provider may decrease, temporarily hold, or permanently stop your treatment with TIBSOVO if you develop certain side effects.

TIBSOVO may cause fertility problems in females and males, which may affect your ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TIBSOVO.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see accompanying Full Prescribing Information, including Medication Guide.

 **TIBSOVO®**
ivosidenib tablets 250mg

HOW SHOULD I STORE TIBSOVO®?

Store TIBSOVO at room temperature from 68 to 77 °F (20 to 25 °C).
Keep TIBSOVO and all medicines out of the reach of children.

GENERAL INFORMATION ABOUT THE SAFE AND EFFECTIVE USE OF TIBSOVO

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not take TIBSOVO for conditions for which it was not prescribed. Do not give TIBSOVO to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about TIBSOVO that is written for healthcare professionals.

WHAT ARE THE INGREDIENTS IN TIBSOVO?

Active ingredient: ivosidenib

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, hypromellose acetate succinate, magnesium stearate, microcrystalline cellulose, and sodium lauryl sulfate. The tablet coating includes FD&C blue #2, hypromellose, lactose monohydrate, titanium dioxide, and triacetin.

For more information, go to www.Tibsovo.com or call 1-800-807-6124.

Please see accompanying Full Prescribing Information, including Medication Guide.

Healthcare team contact information

Name	Name
Phone	Phone
Address	Address
Email	Email

NOTES

NEED HELP WITH ACCESS OR FINANCIAL ASSISTANCE FOR TIBSOVO®?

Talk to your healthcare provider about the **ServierONE Patient Services program** to learn more.

Providers and patients can visit ServierONE.com or call 1-800-813-5905.

For the TIBSOVO copay program, patients can also visit ServierONE-copay.com.

Please see accompanying Full Prescribing Information, including Medication Guide.



© 2021 Servier Pharmaceuticals LLC. Boston, MA 02210. Customer Service: 1-800-807-6124. Servier and the Servier Logo are registered trademarks of LES LABORATOIRES SERVIER. TIBSOVO is a registered trademark of SERVIER PHARMACEUTICALS LLC, a wholly owned, indirect subsidiary of LES LABORATOIRES SERVIER. US-00910 08/2021

