Differentiation Syndrome Reference Card



Remember to carry this card with you at all times.

Scan here to save a copy to your phone.



© 2022 Servier Pharmaceuticals LLC. Boston, MA 02210. Customer Service: 1-800-807-6124. Servier and the Servier Logo are registered trademarks of LES LABORATOIRES SERVIER.

US-01327 v3 07/2022

Information for Patients

There are serious risks associated with your acute myeloid leukemia (AML) therapy, including a condition called differentiation syndrome which may be life-threatening or lead to death.

If you have any of the symptoms to the right while taking treatment, you should call your doctor and seek emergency medical care right away.

These symptoms include:

- Fever
- Cough
- Trouble breathing
- Rash
- Decreased urination
- Dizziness or lightheadedness
- Rapid weight gain
- Swelling of arms or legs

Please see accompanying Full Prescribing Information, including BOXED WARNING.

REMEMBER!



Call your doctor and get emergency medical care right away if you have any of these symptoms. Show this card in the emergency room and to any healthcare provider involved in your care.

Patient name

Prescriber name

Prescriber phone number

Current medication(s)

Information for Healthcare Providers



This patient is taking a medication for the treatment of acute myeloid leukemia (AML) that can cause differentiation syndrome.

Differentiation syndrome may be life-threatening or fatal.

If this patient presents with signs or symptoms of differentiation syndrome:

- Evaluate urgently
- Provide all supportive care
- Contact the prescribing physician (see information at left) as soon as possible to coordinate care

If differentiation syndrome is suspected:

- Administer systemic corticosteroids and initiate hemodynamic monitoring until symptom resolution and for a minimum of 3 days
- Interrupt the AML medication if severe signs and/or symptoms persist for more than 48 hours after initiation of systemic corticosteroids
- Resume the AML medication when signs and symptoms improve to Grade 2 or lower

For additional management information, visit **understand-ds.com**.

To report adverse events, call Servier Pharmaceuticals LLC at 1-800-807-6124.